

The N/O abundance ratio in the lowest-metallicity blue compact dwarf galaxies

Yuri Izotov¹, Frederic Chaffee², Craig Foltz³, Klaus Fricke⁴, Richard Green⁵, Natalia Guseva¹, Kai Noeske⁴, Polychronis Papaderos⁴, Trinh Thuan⁶

¹*Main Astronomical Observatory, 27 Zabolotnoho str., 03680 Kyiv, Ukraine.*

²*W. M. Keck Observatory, 65-1120 Mamalahoa Hwy., Kamuela, HI 96743, USA.*

³*MMT Observatory, University of Arizona, Tucson, AZ 85721, USA.*

⁴*Universitäts-Sternwarte, Geismarlandstraße 11, D-37083 Göttingen, Germany.*

⁵*National Optical Astronomy Observatories, Tucson, AZ 85726, USA.*

⁶*Astronomy Department, University of Virginia, Charlottesville, VA 22903, USA.*

Abstract. The results of the N/O abundance determination in a sample of low-metallicity blue compact dwarf (BCD) galaxies based on new spectroscopic observations with large telescopes (Keck, VLT, MMT, 4m KPNO) are presented. We show that the N/O abundance ratio is constant at lowest metallicities $\leq Z_{\odot}/20$, confirming previous findings and strongly supporting the origin of nitrogen as a primary element.

1 Sample

The sample consists of ~ 100 BCDs with a heavy element abundances in the range $Z_{\odot}/50 - Z_{\odot}/4$, selected from the First and Second Byurakan, Michigan, Hamburg, Tololo and Case surveys. All most metal-deficient BCDs with $Z \leq Z_{\odot}/20$ are included. High signal-to-noise ratio spectroscopic observations were made during 1993 – 2000 with Keck, VLT, MMT and 4m KPNO telescopes. In total, the results of 119 independent observations of H II regions are presented in this contribution.

2 Results

The dependence of the [N/O] vs [O/H] abundance ratios is shown in Fig. 1, where $[N/O] = \log(N/O) - \log(N/O)_{\odot}$ and $[O/H] = \log(O/H) - \log(O/H)_{\odot}$. The open circles show data from [2] and solid circles are new data. We conclude from Fig. 1 that

- the N/O abundance ratio is constant in BCDs with $[O/H] < -1$ and favors the primary production of nitrogen by intermediate-mass and/or massive stars. Secondary nitrogen production is important for higher metallicities;




Figure 1: $[N/O]$ vs $[O/H]$ diagram for BCDs. Open circles are from [2], filled circles are new data. Dotted lines are diagrams for primary and secondary productions of nitrogen, while the solid line shows the dependence $[N/O]$ vs $[O/H]$ when both primary and secondary productions of nitrogen are taken into account.

- the very small dispersion of the N/O abundance ratio in BCDs with $[O/H] < -1.4$ favors the primary production of nitrogen in massive stars;
- the apparent difference of the N/O abundance ratios in BCDs and damped Ly α absorption (DLA) systems (e.g. [1, 2]) could be caused by the different enrichment scenarios or unaccounted ionization effects in DLAs.

Acknowledgements. Y.I. and N.G. have been partially supported by INTAS 97-0033 and Swiss SCOPE 7UKPJ62178 grants. Those authors, P.P. and K.F. acknowledge support by the Volkswagen Foundation under grant No. I/72919. Y.I. and T.T. have been partially supported by NSF grant AST-9616863. Research by P.P. and K.F. has been supported by the Deutsches Zentrum für Luft- und Raumfahrt e.V. (DLR) under grant 50 OR 9907 7. C.F. acknowledges the support of the NSF under grant AST-9803072, and K.N. of the Deutsche Forschungsgemeinschaft (DFG) grant FR325/50-1.

References

- [1] Izotov Y. I., Schaerer D., Charbonnel C., 2000, ApJ 549, 878
- [2] Izotov Y. I., Thuan T. X., 1999, ApJ 511, 639